

1) Find

$$\int_1^9 \frac{4}{\sqrt{x}} dx$$

- a) 71
 - b) 8
 - c) 16
 - d) 24
 - e) 72
- 2) If

$$f'(x) = -4(x-5)^2(x-7)$$

which of the following is true about $y = f(x)$?

- a) f has a local minimum at $x = 5$ and a local maximum at $x = 7$.
 - b) f has a local maximum at $x = 5$ and a local minimum at $x = 7$.
 - c) f has a point of inflection at $x = 5$ and a local maximum at $x = 7$.
 - d) f has a point of inflection at $x = 5$ and a local minimum at $x = 7$.
 - e) f has a local minimum at $x = 5$ and a point of inflection at $x = 7$.
- 3) A curve is described by parametric equations

$$[x = 7 \ln(t), y = t^2 - 5]$$

where $t > 0$. Give an expression for

$$\frac{\partial^2}{\partial x^2} y$$

- a) $\frac{2}{7} t$
- b) $\frac{4}{49} t$
- c) $\frac{2}{7} t^2$
- d) $\frac{4}{49} t^2$
- e) $\frac{4}{7} t$

4) Give the value for

$$\lim_{x \rightarrow 0} \left(\frac{1}{8} \frac{4^x - 1}{x} \right)$$

- a) *The limit does not exist.*
- b) 1
- c) $16 \ln(2)$
- d) $\frac{1}{4} \ln(2)$
- e) 0

5) Which of the following series converge?

I.

$$\sum_{n=1}^{\infty} \frac{\sin(n)}{n!}$$

II.

$$\sum_{n=1}^{\infty} \frac{x^{2n}}{n!}$$

III.

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)}$$

a) II and III only

b) II only

c) I, II and III

d) I and II only

e) I only

6) If $g(f(x)) = x$, $g(5) = 2$ and $g'(5) = 12$, then $f'(2)$ is

a) $-\frac{1}{12}$

b) $\frac{1}{12}$

c) $\frac{1}{5}$

d) $-\frac{1}{5}$

e) $\frac{5}{12}$

7) If f is a differentiable function and $f(0) = -6$ and $f(4) = 12$, then which of the following must be true?

I. There exists a c in $[0,4]$ where $f(c) = 0$.

II. There exists a c in $[0,4]$ where $f'(c) = 0$.

III. There exists a c in $[0,4]$ where $f'(c) = 9/2$.

a) I and III only

b) I only

c) II only

d) II and III only

e) I, II and III

8) Evaluate

$$\int_{-2}^2 \frac{7}{x^2} dx$$

a) 0

b) *The integral diverges.*

c) $\frac{7}{12}$

d) $\frac{7}{6}$

e) $\frac{7}{24}$

9) Find the area enclosed by the graphs of

$$\begin{aligned}y &= e^x + 1 \\ y &= 5\end{aligned}$$

and the y-axis.

a) $5 \ln(5) - 5$

b) $8 \ln(2) - 3$

c) $5 \ln(5) + 5$

d) $10 \ln(2) + 4$

e) $10 \ln(2) - 4$

10) What is the minimum value of the function

$$f(x) = \frac{3}{\sqrt{x}} + 6\sqrt{x}$$

a) $\frac{1}{2}$

b) $3\sqrt{2}$

c) $6\sqrt{2}$

d) $\frac{1}{2}\sqrt{2}$

e) $\sqrt{2}$

11) Give the value of

$$\int_{\pi}^{2\pi} \frac{\cos(7x)}{2 + \sin(7x)} dx$$

a) 0

b) 1

c) $\frac{1}{7}$

d) $\frac{1}{14}$

e) $-\frac{1}{7}$

12) The side of a cube is expanding at a constant rate of 3 inches per second. What is the rate of change of the surface area, in in^2 per second, when the volume of the cube is 64 in^3 ?

a) 72

b) 36

c) 144

d) 180

e) 18

13) Give the area inside one petal of the polar graph of

$$r = 4 \sin(2\theta)$$

- a) 2π
- b) 4π
- c) 8π
- d) π
- e) 2

14) Give the solution to the initial value problem

$$[y' = 12x^2y, y(1) = 1]$$

- a) $\frac{1}{4} \ln(x^3) + e$
- b) e^{4x^3}
- c) $e^{4x^3} - 4$
- d) $\frac{1}{4} \ln(x^3) + 1$
- e) $e^{4x^3} - 4$

15) The position of a particle moving along a horizontal line is given by

$$x(t) = 2(t-3)^3$$

What is the maximum speed of the particle for $0 \leq t \leq 10$?

- a) 18
- b) 54
- c) 162
- d) 294
- e) 98

16) $\int [\sec(7x)]^2 dx =$

- a) $-7 \tan(7x) + C$
- b) $7 \tan(7x) + C$
- c) $\frac{1}{7} \tan(7x) + C$
- d) $7 [\tan(7x)]^2 + C$
- e) $\frac{1}{7} [\tan(7x)]^2 + C$


17) Define the function

$$f(x) = x e^{-2x}$$

for $x > 0$. Give the interval on which the function is increasing.

- a) $\left(0, \frac{1}{2}\right)$
- b) $(1, 2)$
- c) $\left(1, \frac{1}{2}e\right)$
- d) $\left(0, \frac{1}{2}e\right)$
- e) $(0, 2)$

18)



Which of the following differential equations correspond to the slope field shown in the figure above?

- a) $\frac{dy}{dx} = \frac{x}{y}$
- b) $\frac{dy}{dx} = \frac{y}{x}$
- c) $\frac{dy}{dx} = -\frac{y}{x}$
- d) $\frac{dy}{dx} = 6xy$
- e) $\frac{dy}{dx} = -\frac{x}{y}$

19) Evaluate

$$\lim_{h \rightarrow 0} \left(\frac{\cos(2x + 2h) - \cos(2x)}{h} \right)$$

- a) $2 \sin(2x)$
- b) $-2 \sin(2x)$
- c) $2 \cos(2x)$
- d) $-2 \cos(2x)$
- e) *The limit does not exist.*

20) If $\int_0^{25} e^x dx = m$ then $\int_0^5 x e^{x^2} dx$ is

- a) m^2
- b) m
- c) $2m$
- d) $\frac{1}{2} m^2$
- e) $\frac{1}{2} m$

21) Find the area of the region enclosed by the graph of

$$y = 4x^2$$

and the line

$$y = 2x$$

- a) $\frac{1}{12}$
- b) $\frac{1}{6}$
- c) $\frac{1}{24}$
- d) $\frac{8}{3}$
- e) $\frac{16}{3}$

22) Suppose

$$z = e^y$$

$$y = 2x^3 - 2$$

and

$$x = 1 + \ln(t^2)$$

What is dz/dt when $t = 1$?

- a) 12
- b) 2
- c) 6
- d) 3
- e) 8

23) Evaluate

$$\int_0^{\frac{1}{2}\pi} \cos^2(5x) \sin(5x) \, dx$$

- a) $\frac{1}{5}$
- b) $\frac{1}{15}$
- c) $\frac{1}{30}$
- d) $-\frac{1}{15}$
- e) $-\frac{1}{30}$

24) Give an equation for the tangent line to the parametric curve

$$[x = e^t, y = t^2 + 5t]$$

at $t = 0$.

- a) $y - 1 = 5x$
- b) $y = \frac{5(x-1)}{e}$
- c) $y = 5e(x-1)$
- d) $y - 1 = 5x - 5$
- e) $y = 5x - 5$

25) Evaluate

$$\frac{\partial}{\partial x} \int_4^{8x} \ln(4t) dt$$

- a) $8 \ln(8x) - 32 \ln(2)$
- b) $4 \ln(32x)$
- c) $8 \ln(32x) - 32 \ln(2)$
- d) $8 \ln(32x)$
- e) $\frac{2}{x}$

26) The region bounded by

$$y = 2 \sin(x)$$

and the x -axis, for $0 \leq x \leq \frac{1}{2} \pi$, is rotated about the line $y = -4$. The volume of this solid can be represented by:

- a) $2 \pi \int_0^{\frac{1}{2} \pi} (2 \sin(x) + 4)^2 dx$
- b) $\pi \int_0^{\frac{1}{2} \pi} (4 \sin(x)^2 - 16) dx$
- c) $2 \pi \int_0^{\frac{1}{2} \pi} 4 \sin(x + 4)^2 dx$
- d) $\pi \int_0^{\frac{1}{2} \pi} ((2 \sin(x) + 4)^2 - 16) dx$
- e) $2 \pi \int_0^{\frac{1}{2} \pi} (4 \sin(x)^2 + 4) dx$

27) Give the third degree Taylor polynomial about $x = 1$ of

$$f(x) = \ln(x)$$

- a) $(x-1) + (x-1)^2 + 2(x-1)^3$
- b) $(x-1) - \frac{1}{3}(x-1)^2 + \frac{1}{5}(x-1)^3$
- c) $(x-1) + \frac{1}{3}(x-1)^2 + \frac{1}{3}(x-1)^3$
- d) $(x-1) - \frac{1}{2}(x-1)^2 + \frac{1}{3}(x-1)^3$
- e) $(x-1) - \frac{1}{2}(x-1)^2 + \frac{1}{6}(x-1)^3$

28) Which of the following integrals gives the length of the graph of

$$f(x) = e^{5x}$$

for x between 0 and 2?

a) $\int_0^2 \sqrt{x + 25 e^{10x}} \, dx$

b) $\int_0^2 \sqrt{1 + e^{10x}} \, dx$

c) $\int_0^2 \sqrt{1 + 25 e^{10x}} \, dx$

d) $\int_0^2 \sqrt{x + e^{10x}} \, dx$

e) $\int_0^2 \sqrt{e^{5x} + 25 e^{10x}} \, dx$

29) Find the average value of the function

$$f(x) = e^{7x}$$

over the interval $[0, 4]$.

- a) $\frac{1}{28} e^{28}$
- b) $\frac{1}{7} (e^{28} - 1)$
- c) $\frac{1}{28} (e^{28} - 1)$
- d) $\frac{1}{4} (e^{28} - 1)$
- e) $\frac{1}{7} e^{28}$

30) What is the y-intercept of the line tangent to the curve $y = x^2 + 4$ at $x = 3$?

- a) $(0, -14)$
- b) $(0, -2)$
- c) $(0, -5)$
- d) $(0, 14)$
- e) $(0, 5)$

31) Which of the following function(s) is continuous and differentiable?

I. $f(x) = \frac{7}{\sqrt{x}}$

II. $g(x) = x|x|$

III. $h(x) = \begin{cases} 4x + 1 & x \leq 0 \\ x^2 + 1 & 0 < x \end{cases}$


- a) I and III only
- b) III only
- c) I and II only
- d) I only
- e) II only

32) Find m

$$\lim_{x \rightarrow 0} \left(\frac{e^{mx^2} - \cos(6x)}{x^2} \right) = 36$$

- a) $\frac{3}{2}$
- b) 3
- c) 9
- d) 18
- e) 72

33) The graph of the derivative of f is shown below. Which of the following must be true?



a) f has a local maximum at $x = 0$.

b) f is increasing on $[-6, 6]$.

c) f has a point of inflection at $x = 12$.

d) f is concave down on $[0, 12]$.

e) f has a local minimum at $x = -6$.

34) The sum of two positive integers x and y is 90. Find the value of x that minimizes

$$P = x^3 - 90xy$$

a) $x = 45$

b) $x = 30$

c) $x = 15$

d) $x = 75$

e) $x = 60$

35) A particle moves on the curve

$$[x = 4 \sin(t), y = \sin(2t)]$$

find the speed of the particle at time $t = \pi$.

a) 4.4721

b) 3.4641

c) 4.2426

d) 2.8284

e) 2.4495

36) The function f is defined as

$$f(x) = \frac{(x-2)^2}{x-7}$$

$$x \neq 7$$

Which of the following is **false**?

a) f has a vertical asymptote at $x = 7$.

b) f has a horizontal asymptote at $y = 1$.

c) f is decreasing on $[2, 7]$.

d) f has a local maximum at $x = 2$.

e) f is concave up for $x > 7$.

37) A particle is moving along the x -axis and its position at time $t \geq 0$ is given by

$$S(t) = (t-2)^2(t-5)$$

Which of the following is (are) true?

I. The particle changes direction at $x = 2$ and $x = 5$.

II. The particle is slowing down on $[0, 2]$.

III. The particle is speeding up on $[2, 5]$.

- a) I only
- b) II only
- c) II and III only
- d) I and III only
- e) I, II and III

38) $f(x)$ is a differentiable function and it is decreasing on $(-\infty, \infty)$.

If

$$g(x) = f(8x^3 - x^2)$$

then g has a local maximum at

- a) $x = 1$
- b) $x = 0$
- c) $x = \frac{1}{12}$
- d) $x = 4$
- e) $x = \frac{1}{4}$


39) The rate at which a bacteria population grows is proportional to the number of bacteria present. Initially, there were 1000 bacteria present and the population doubled in 3 hours. Roughly how many hours does it take for the population to reach 10000?

- a) 9.9
- b) 12.3
- c) 15.6
- d) 6.6
- e) 8.7

40) Find the radius of convergence of the series

$$\sum_{n=1}^{\infty} \frac{n(5x-11)^n}{2^n}$$

- a) $\frac{2}{11}$
- b) $\frac{5}{2}$
- c) $\frac{2}{5}$
- d) $\frac{11}{5}$
- e) $\frac{11}{2}$



41)

$$g(x) = \int_0^x f(t) dt$$

for $0 \leq x \leq 9$

The graph of g is shown above. Which of the following must be true?

- I. $\int_0^9 f(t) dt = 0$
- II. $\int_3^6 f(t) dt = 3$
- III. $\int_6^0 f(t) dt = -3$

- a) I and II only
- b) I only
- c) II only
- d) I and III only
- e) II and III only

42) If the region bounded by $y = \tan^{-1}(x)$, $y = \frac{1}{4}\pi$ and the y -axis is rotated about the y -axis, the volume of

the solid formed is

- a) 0.430
- b) 0.215
- c) 1.348
- d) 0.413
- e) 0.674

43) $f(x)$ is represented by the Maclaurin series

$$1 - \frac{(2x)^2}{2!} + \frac{(2x)^4}{4!} - \dots + (-1)^n \frac{(2x)^{2n}}{(2n)!} + \dots$$

What is the slope of the line normal to the graph of f at $x = \frac{11}{4}\pi$?

- a) -2
- b) 0
- c) 2
- d) $-\frac{1}{2}$

e) $\frac{1}{2}$

44) What are all values of h for which

$$\int_0^{\infty} \frac{8x}{(x^2 + 1)^h} dx$$

converge?

a) $h \geq 1$

b) $h < 1$

c) $h > 1$

d) $h \leq 1$

e) $-1 < h < 1$

45) The base of a solid is the region bounded by

$$y = 2\sqrt{x}$$

the x -axis, and

the line $x = 2$

Each cross-section of the solid perpendicular to the x -axis is a square, with one side on the xy -plane. Which of the following expressions represents the volume of the solid?

a) $\int_0^2 4x \, dx$

b) $\int_0^2 2\sqrt{x} \, dx$

c) $\int_0^2 2x \, dx$

d) $\int_0^1 4x \, dx$

e) $\int_0^1 2\sqrt{x} \, dx$

1) Find

$$\int_1^9 \frac{4}{\sqrt{x}} dx$$

- a) 71
 - b) 8
 - *c) 16
 - d) 24
 - e) 72
- 2) If

$$f'(x) = -4(x-5)^2(x-7)$$

which of the following is true about $y = f(x)$?

- a) f has a local minimum at $x = 5$ and a local maximum at $x = 7$.
 - b) f has a local maximum at $x = 5$ and a local minimum at $x = 7$.
 - *c) f has a point of inflection at $x = 5$ and a local maximum at $x = 7$.
 - d) f has a point of inflection at $x = 5$ and a local minimum at $x = 7$.
 - e) f has a local minimum at $x = 5$ and a point of inflection at $x = 7$.
- 3) A curve is described by parametric equations

$$[x = 7 \ln(t), y = t^2 - 5]$$

where $t > 0$. Give an expression for

$$\frac{\partial^2}{\partial x^2} y$$

- a) $\frac{2}{7} t$
- b) $\frac{4}{49} t$
- c) $\frac{2}{7} t^2$
- *d) $\frac{4}{49} t^2$
- e) $\frac{4}{7} t$

4) Give the value for

$$\lim_{x \rightarrow 0} \left(\frac{1}{8} \frac{4^x - 1}{x} \right)$$

- a) *The limit does not exist.*
- b) 1
- c) $16 \ln(2)$
- *d) $\frac{1}{4} \ln(2)$
- e) 0

5) Which of the following series converge?

I.

$$\sum_{n=1}^{\infty} \frac{\sin(n)}{n!}$$

II.

$$\sum_{n=1}^{\infty} \frac{x^{2n}}{n!}$$

III.

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)}$$

a) II and III only

b) II only

*c) I, II and III

d) I and II only

e) I only

6) If $g(f(x)) = x$, $g(5) = 2$ and $g'(5) = 12$, then $f'(2)$ is

a) $-\frac{1}{12}$

*b) $\frac{1}{12}$

c) $\frac{1}{5}$

d) $-\frac{1}{5}$

e) $\frac{5}{12}$

7) If f is a differentiable function and $f(0) = -6$ and $f(4) = 12$, then which of the following must be true?

I. There exists a c in $[0,4]$ where $f(c) = 0$.

II. There exists a c in $[0,4]$ where $f'(c) = 0$.

III. There exists a c in $[0,4]$ where $f'(c) = 9/2$.

*a) I and III only

b) I only

c) II only

d) II and III only

e) I, II and III

8) Evaluate

$$\int_{-2}^2 \frac{7}{x^2} dx$$

a) 0

*b) *The integral diverges.*

c) $\frac{7}{12}$

d) $\frac{7}{6}$

e) $\frac{7}{24}$

9) Find the area enclosed by the graphs of

$$\begin{aligned}y &= e^x + 1 \\ y &= 5\end{aligned}$$

and the y-axis.

a) $5 \ln(5) - 5$

*b) $8 \ln(2) - 3$

c) $5 \ln(5) + 5$

d) $10 \ln(2) + 4$

e) $10 \ln(2) - 4$

10) What is the minimum value of the function

$$f(x) = \frac{3}{\sqrt{x}} + 6\sqrt{x}$$

a) $\frac{1}{2}$

b) $3\sqrt{2}$

*c) $6\sqrt{2}$

d) $\frac{1}{2}\sqrt{2}$

e) $\sqrt{2}$

11) Give the value of

$$\int_{\pi}^{2\pi} \frac{\cos(7x)}{2 + \sin(7x)} dx$$

*a) 0

b) 1

c) $\frac{1}{7}$

d) $\frac{1}{14}$

e) $-\frac{1}{7}$

12) The side of a cube is expanding at a constant rate of 3 inches per second. What is the rate of change of the surface area, in in^2 per second, when the volume of the cube is 64 in^3 ?

a) 72

b) 36

*c) 144

d) 180

e) 18

13) Give the area inside one petal of the polar graph of

$$r = 4 \sin(2\theta)$$

- *a) 2π
- b) 4π
- c) 8π
- d) π
- e) 2

14) Give the solution to the initial value problem

$$[y' = 12x^2y, y(1) = 1]$$

- a) $\frac{1}{4} \ln(x^3) + e$
- b) e^{4x^3}
- c) $e^{4x^3} - 4$
- d) $\frac{1}{4} \ln(x^3) + 1$
- *e) $e^{4x^3} - 4$

15) The position of a particle moving along a horizontal line is given by

$$x(t) = 2(t-3)^3$$

What is the maximum speed of the particle for $0 \leq t \leq 10$?

- a) 18
- b) 54
- c) 162
- *d) 294
- e) 98

16) $\int [\sec(7x)]^2 dx =$

- a) $-7 \tan(7x) + C$
- b) $7 \tan(7x) + C$
- *c) $\frac{1}{7} \tan(7x) + C$
- d) $7 [\tan(7x)]^2 + C$
- e) $\frac{1}{7} [\tan(7x)]^2 + C$


17) Define the function

$$f(x) = x e^{-2x}$$

for $x > 0$. Give the interval on which the function is increasing.

- *a) $\left(0, \frac{1}{2}\right)$
- b) $(1, 2)$
- c) $\left(1, \frac{1}{2}e\right)$
- d) $\left(0, \frac{1}{2}e\right)$
- e) $(0, 2)$

18)



Which of the following differential equations correspond to the slope field shown in the figure above?

- a) $\frac{dy}{dx} = \frac{x}{y}$
- b) $\frac{dy}{dx} = \frac{y}{x}$
- c) $\frac{dy}{dx} = -\frac{y}{x}$
- d) $\frac{dy}{dx} = 6xy$
- *e) $\frac{dy}{dx} = -\frac{x}{y}$

19) Evaluate

$$\lim_{h \rightarrow 0} \left(\frac{\cos(2x + 2h) - \cos(2x)}{h} \right)$$

- a) $2 \sin(2x)$
- *b) $-2 \sin(2x)$
- c) $2 \cos(2x)$
- d) $-2 \cos(2x)$
- e) *The limit does not exist.*

20) If $\int_0^{25} e^x dx = m$ then $\int_0^5 x e^{x^2} dx$ is

- a) m^2
- b) m
- c) $2m$
- d) $\frac{1}{2} m^2$
- *e) $\frac{1}{2} m$

21) Find the area of the region enclosed by the graph of

$$y = 4x^2$$

and the line

$$y = 2x$$

*a) $\frac{1}{12}$

b) $\frac{1}{6}$

c) $\frac{1}{24}$

d) $\frac{8}{3}$

e) $\frac{16}{3}$

22) Suppose

$$\begin{aligned}z &= e^y \\ y &= 2x^3 - 2\end{aligned}$$

and

$$x = 1 + \ln(t^2)$$

What is dz/dt when $t = 1$?

*a) 12

b) 2

c) 6

d) 3

e) 8

23) Evaluate

$$\int_0^{\frac{1}{2}\pi} \cos^2(5x) \sin(5x) \, dx$$

a) $\frac{1}{5}$

*b) $\frac{1}{15}$

c) $\frac{1}{30}$

d) $-\frac{1}{15}$

e) $-\frac{1}{30}$

24) Give an equation for the tangent line to the parametric curve

$$[x = e^t, y = t^2 + 5t]$$

at $t = 0$.

a) $y - 1 = 5x$

b) $y = \frac{5(x-1)}{e}$

c) $y = 5e(x-1)$

d) $y - 1 = 5x - 5$

*e) $y = 5x - 5$

25) Evaluate

$$\frac{\partial}{\partial x} \int_4^{8x} \ln(4t) dt$$

- a) $8 \ln(8x) - 32 \ln(2)$
- b) $4 \ln(32x)$
- c) $8 \ln(32x) - 32 \ln(2)$
- *d) $8 \ln(32x)$
- e) $\frac{2}{x}$

26) The region bounded by

$$y = 2 \sin(x)$$

and the x -axis, for $0 \leq x \leq \frac{1}{2} \pi$, is rotated about the line $y = -4$. The volume of this solid can be represented by:

- a) $2\pi \int_0^{\frac{1}{2}\pi} (2 \sin(x) + 4)^2 dx$
- b) $\pi \int_0^{\frac{1}{2}\pi} (4 \sin(x)^2 - 16) dx$
- c) $2\pi \int_0^{\frac{1}{2}\pi} 4 \sin(x + 4)^2 dx$
- *d) $\pi \int_0^{\frac{1}{2}\pi} ((2 \sin(x) + 4)^2 - 16) dx$
- e) $2\pi \int_0^{\frac{1}{2}\pi} (4 \sin(x)^2 + 4) dx$

27) Give the third degree Taylor polynomial about $x = 1$ of

$$f(x) = \ln(x)$$

- a) $(x-1) + (x-1)^2 + 2(x-1)^3$
- b) $(x-1) - \frac{1}{3}(x-1)^2 + \frac{1}{5}(x-1)^3$
- c) $(x-1) + \frac{1}{3}(x-1)^2 + \frac{1}{3}(x-1)^3$
- *d) $(x-1) - \frac{1}{2}(x-1)^2 + \frac{1}{3}(x-1)^3$
- e) $(x-1) - \frac{1}{2}(x-1)^2 + \frac{1}{6}(x-1)^3$

28) Which of the following integrals gives the length of the graph of

$$f(x) = e^{5x}$$

for x between 0 and 2?

a) $\int_0^2 \sqrt{x + 25 e^{10x}} dx$

b) $\int_0^2 \sqrt{1 + e^{10x}} dx$

*c) $\int_0^2 \sqrt{1 + 25 e^{10x}} dx$

d) $\int_0^2 \sqrt{x + e^{10x}} dx$

e) $\int_0^2 \sqrt{e^{5x} + 25 e^{10x}} dx$

29) Find the average value of the function

$$f(x) = e^{7x}$$

over the interval $[0, 4]$.

- a) $\frac{1}{28} e^{28}$
- b) $\frac{1}{7} (e^{28} - 1)$
- *c) $\frac{1}{28} (e^{28} - 1)$
- d) $\frac{1}{4} (e^{28} - 1)$
- e) $\frac{1}{7} e^{28}$

30) What is the y-intercept of the line tangent to the curve $y = x^2 + 4$ at $x = 3$?

- a) $(0, -14)$
- b) $(0, -2)$
- *c) $(0, -5)$
- d) $(0, 14)$
- e) $(0, 5)$

31) Which of the following function(s) is continuous and differentiable?

I. $f(x) = \frac{7}{\sqrt{x}}$

II. $g(x) = x|x|$

III. $h(x) = \begin{cases} 4x + 1 & x \leq 0 \\ x^2 + 1 & 0 < x \end{cases}$


- a) I and III only
- b) III only
- c) I and II only
- *d) I only
- e) II only

32) Find m

$$\lim_{x \rightarrow 0} \left(\frac{e^{mx^2} - \cos(6x)}{x^2} \right) = 36$$

- a) $\frac{3}{2}$
- b) 3
- c) 9
- *d) 18
- e) 72

33) The graph of the derivative of f is shown below. Which of the following must be true?



a) f has a local maximum at $x = 0$.

b) f is increasing on $[-6, 6]$.

*c) f has a point of inflection at $x = 12$.

d) f is concave down on $[0, 12]$.

e) f has a local minimum at $x = -6$.

34) The sum of two positive integers x and y is 90. Find the value of x that minimizes

$$P = x^3 - 90xy$$

a) $x = 45$

*b) $x = 30$

c) $x = 15$

d) $x = 75$

e) $x = 60$

35) A particle moves on the curve

$$[x = 4 \sin(t), y = \sin(2t)]$$

find the speed of the particle at time $t = \pi$.

*a) 4.4721

b) 3.4641

c) 4.2426

d) 2.8284

e) 2.4495

36) The function f is defined as

$$f(x) = \frac{(x-2)^2}{x-7}$$

$$x \neq 7$$

Which of the following is **false**?

a) f has a vertical asymptote at $x = 7$.

*b) f has a horizontal asymptote at $y = 1$.

c) f is decreasing on $[2, 7]$.

d) f has a local maximum at $x = 2$.

e) f is concave up for $x > 7$.

37) A particle is moving along the x -axis and its position at time $t \geq 0$ is given by

$$S(t) = (t-2)^2(t-5)$$

Which of the following is (are) true?

I. The particle changes direction at $x = 2$ and $x = 5$.

II. The particle is slowing down on $[0, 2]$.

III. The particle is speeding up on $[2, 5]$.

a) I only

*b) II only

c) II and III only

d) I and III only

e) I, II and III

38) $f(x)$ is a differentiable function and it is decreasing on $(-\infty, \infty)$.

If

$$g(x) = f(8x^3 - x^2)$$

then g has a local maximum at

a) $x = 1$

b) $x = 0$

*c) $x = \frac{1}{12}$

d) $x = 4$

e) $x = \frac{1}{4}$

39) The rate at which a bacteria population grows is proportional to the number of bacteria present. Initially, there were 1000 bacteria present and the population doubled in 3 hours. Roughly how many hours does it take for the population to reach 10000?

*a) 9.9

b) 12.3

c) 15.6

d) 6.6

e) 8.7

40) Find the radius of convergence of the series

$$\sum_{n=1}^{\infty} \frac{n(5x-11)^n}{2^n}$$


a) $\frac{2}{11}$

b) $\frac{5}{2}$

*c) $\frac{2}{5}$

d) $\frac{11}{5}$

e) $\frac{11}{2}$



41)

$$g(x) = \int_0^x f(t) dt$$

for $0 \leq x \leq 9$

The graph of g is shown above. Which of the following must be true?

- I. $\int_0^9 f(t) dt = 0$
- II. $\int_3^6 f(t) dt = 3$
- III. $\int_6^0 f(t) dt = -3$

- *a) I and II only
- b) I only
- c) II only
- d) I and III only
- e) II and III only

42) If the region bounded by $y = \tan^{-1}(x)$, $y = \frac{1}{4} \pi$ and the y-axis is rotated about the y-axis, the volume of

the solid formed is

- a) 0.430
- b) 0.215
- c) 1.348
- d) 0.413
- *e) 0.674

43) $f(x)$ is represented by the Maclaurin series

$$1 - \frac{(2x)^2}{2!} + \frac{(2x)^4}{4!} - \dots + (-1)^n \frac{(2x)^{2n}}{(2n)!} + \dots$$

What is the slope of the line normal to the graph of f at $x = \frac{11}{4} \pi$?

- a) -2
- b) 0
- c) 2
- *d) $-\frac{1}{2}$

e) $\frac{1}{2}$

44) What are all values of h for which

$$\int_0^{\infty} \frac{8x}{(x^2 + 1)^h} dx$$

converge?

a) $h \geq 1$

b) $h < 1$

*c) $h > 1$

d) $h \leq 1$

e) $-1 < h < 1$

45) The base of a solid is the region bounded by

$$y = 2\sqrt{x}$$

the x -axis, and

the line $x = 2$

Each cross-section of the solid perpendicular to the x -axis is a square, with one side on the xy -plane. Which of the following expressions represents the volume of the solid?

*a) $\int_0^2 4x dx$

b) $\int_0^2 2\sqrt{x} dx$

c) $\int_0^2 2x dx$

d) $\int_0^1 4x dx$

e) $\int_0^1 2\sqrt{x} dx$