1) Find

$$\int_{1}^{9} \frac{2}{\sqrt{x}} \, \mathrm{d}x$$

- a) ₁₂
- b) 4
- c) 35
- d) ₃₆
- e) 8
- 2) If

$$f'(x) = -5 (x-6)^2 (x-9)$$

which of the following is true about y = f(x)? a) *f* has a local maximum at x = 6 and a local minimum at x = 9. b) *f* has a point of inflection at x = 6 and a local maximum at x = 9. c) *f* has a local minimum at x = 6 and a local maximum at x = 9. d) *f* has a point of inflection at x = 6 and a local minimum at x = 9. e) *f* has a local minimum at x = 6 and a local minimum at x = 9. f has a local minimum at x = 6 and a local minimum at x = 9. f has a local minimum at x = 6 and a local minimum at x = 9. a) *f* has a local minimum at x = 6 and a point of inflection at x = 9. b) *f* has a local minimum at x = 6 and a point of inflection at x = 9. c) *f* has a local minimum at x = 6 and a point of inflection at x = 9. c) *f* has a local minimum at x = 6 and a point of inflection at x = 9. c) *f* has a local minimum at x = 6 and a point of inflection at x = 9. c) *f* has a local minimum at x = 6 and a point of inflection at x = 9. c) *f* has a local minimum at x = 6 and a point of inflection at x = 9. c) *f* has a local minimum at x = 6 and a point of inflection at x = 9. c) *f* has a local minimum at x = 6 and a point of inflection at x = 9. c) *f* has a local minimum at x = 6 and a point of inflection at x = 9. c) *f* has a local minimum at x = 6 and a point of inflection at x = 9. c) *f* has a local minimum at x = 6 and a point of inflection at x = 9. c) *f* has a local minimum at x = 6 and a point of inflection at x = 9. c) *f* has a local minimum at x = 6 and a point of inflection at x = 9. c) *f* has a local minimum at x = 6 and a point of inflection at x = 9. c) *f* has a local minimum at x = 6 and a point of inflection at x = 9. c) *f* has a local minimum at x = 6 and x = 6 and x = 6. c) *f* has a local minimum at x = 6. c) *f* has a local minimum at x = 6. c) *f* has a local minimum at x = 6. c) *f* has a local minimum at x = 6. c) *f* has a local minimum at x = 6.

$$\left[x=5\ln(t),\,y=t^2-4\right]$$

where t > 0. Give an expression for

$$\frac{\partial^2}{\partial x^2}y$$

a) $\frac{4}{25}t^2$ b) $\frac{4}{5}t$ c) $\frac{4}{25}t$ d) $\frac{2}{5}t^2$ e) $\frac{2}{5}t$

4) Give the value for

$$\lim_{x\to 0} \left(\frac{1}{7} \ \frac{5^x - 1}{x}\right)$$

- a) $7 \ln(5)$
- b) ₍₎
- c) 1

d) The limit does not exist.

e)
$$\frac{1}{7} \ln(5)$$

5) Which of the following series converge?

I.

$$\sum_{n=1}^{\infty} \frac{\sin(n)}{n!}$$

II.

$$\sum_{n=1}^{\infty} \frac{x^{2n}}{n!}$$

III.

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)}$$

a) I, II and III b) II only c) II and III only d) I and II only e) I only 6) If g(f(x)) = x, g(4) = 2 and g'(4) = 10, then f'(2) is a) $\frac{1}{4}$ b) $-\frac{1}{10}$ c) $\frac{1}{10}$ d) $-\frac{1}{4}$ e) $\frac{2}{5}$

7) If *f* is a differentiable function and f(0) = -5 and f(3) = 10, then which of the following must be true?

I. There exists a *c* in [0,3] where f(c) = 0. II. There exists a *c* in [0,3] where f'(c) = 0. III. There exists a *c* in [0,3] where f'(c) = 5.

a) I only

- b) I and III only
- c) II only
- d) II and III only
- e) I, II and III
- 8) Evaluate



a) <u>8</u> 81

- b) ₀
- c) The integral diverges.

d) $\frac{16}{81}$ e) $\frac{4}{81}$

9) Find the area enclosed by the graphs of

$$y = e^{x} + 1$$
$$y = 6$$

and the y-axis.

- a) $6\ln(5) 5$
- b) $6\ln(6) + 6$
- c) $6\ln(5) + 5$
- d) $5\ln(5) 4$
- e) $6\ln(6) 6$

10) What is the minimum value of the function

$$f(x) = \frac{2}{\sqrt{x}} + 3\sqrt{x}$$

a) $\frac{2}{3}\sqrt{6}$ b) $\frac{2}{3}$ c) $\frac{1}{3}\sqrt{6}$ d) $2\sqrt{6}$ e) $\sqrt{6}$ 11) Give the value of

$$\int_{\pi}^{2\pi} \frac{\cos(8x)}{2+\sin(8x)} \, \mathrm{d}x$$

- a) $-\frac{1}{8}$ b) $\frac{1}{8}$ c) $\frac{1}{16}$ d) $_{0}$
- e) 1

12) The side of a cube is expanding at a constant rate of 6 inches per second. What is the rate of change of the surface area, in in^2 per second, when the volume of the cube is 64 in³?

- a) ₃₆
- b) ₁₄₄
- c) ₃₆₀
- d) ₂₈₈
- e) ₇₂
- 13) Give the area inside one petal of the polar graph of

 $r = 5 \sin(2 \theta)$

a) $\frac{25}{2} \pi$ b) $\frac{25}{4} \pi$ c) $\frac{25}{8} \pi$ d) $\frac{25}{16} \pi$ e) $\frac{5}{2}$

14) Give the solution to the initial value problem

$$[y' = 9 x^2 y, y(1) = 1]$$

a) $\frac{1}{3} \ln(x^3) + 1$ b) $e^{3x^3} - 3$ c) $\frac{1}{3} \ln(x^3) + e$ d) $e^{3x^3} - 3$ e) $e^{3x^3} - 3$

15) The position of a particle moving along a horizontal line is given by

 $x(t) = 3 (t-4)^3$ What is the maximum speed of the particle for 0 < t < 10?

- a) 48
- b) ₁₄₄
- c) 576
- d) ₃₂₄
- e) ₁₀₈
- 16) $\int [\sec(3x)]^2 dx =$ a) $\frac{1}{3} [\tan(3x)]^2 + C$
- b) $-3\tan(3x) + C$
- c) $3 [\tan(3x)]^2 + C$ d) $\frac{1}{3} \tan(3x) + C$
- e) $3 \tan(3x) + C$ 17) Define the function

 $f(x) = x e^{-8x}$

for x > 0. Give the interval on which the function is increasing. a) $\left(0, \frac{1}{8}\right)$

b) (1, 8)c) $(1, \frac{1}{8} e)$

d) $\left(0, \frac{1}{8}e\right)$								
e) (0, 8) 18)								
1	1	1	-		1	~	\sim	
1	1	1	-		~	~	N	
7	1	7	1		\$	×.	X.	
T	1	1	1	~	N	1	Ţ	
l	1	1	1	7	1	1	1	
I	X	X	\sim	-	1	1	1	
V.	\mathbf{v}	\sim	-	-	1	1	1	
\mathbf{x}	\sim	~	-		1	1	1	

Which of the following differential equations correspond to the slope field shown in the figure above?

a) $\frac{dy}{dx} = -\frac{x}{y}$ b) $\frac{dy}{dx} = \frac{y}{x}$ c) $\frac{dy}{dx} = -\frac{y}{x}$ d) $\frac{dy}{dx} = \frac{x}{y}$ e) $\frac{dy}{dx} = 5 x y$ 19) Evaluate $\lim_{h \to 0} \left(\frac{\cos(5x+5h) - \cos(5x)}{h} \right)$ a) $-5\sin(5x)$ b) $5\sin(5x)$ c) $5\cos(5x)$ d) $-5\cos(5x)$ e) The limit does not exist. 20) If $\int_{0}^{16} e^{x} dx = m \text{ then } \int_{0}^{4} x e^{x^{2}} dx \text{ is}$ a) $\frac{1}{2}m$ b) _m c) 2 m d) _m2 e) $\frac{1}{2}m^2$ 21) Find the area of the region enclosed by the graph of

and the line

$$y=2x^2$$

$$y = 3x$$

a) $\frac{9}{8}$ b) $\frac{9}{4}$ c) $\frac{3}{8}$ d) $\frac{4}{27}$ e) $\frac{8}{27}$ 22) Suppose

 $z = e^{y}$ $y = 4x^{3} - 4$ $x = 1 + Ln(t^{2})$

and

What is dz/dt when t = 1?

- a) 6
- b) 4
- c) ₁₂
- d) ₁₆
- e) ₂₄
- 23) Evaluate

$$\int_{0}^{\frac{1}{2}\pi} \cos^2\left(13x\right) \sin(13x) \, \mathrm{d}x$$

- a) $-\frac{1}{78}$ b) $\frac{1}{78}$ c) $-\frac{1}{39}$ d) $\frac{1}{39}$
- e) $\frac{1}{13}$

24) Give an equation for the tangent line to the parametric curve $[x = e^t, y = t^2 + 4t]$

at t = 0. a) $y = \frac{4(x-1)}{e}$ b) y = 4x-4c) y = 4e(x-1) d) y-1 = 4xe) y-1 = 4x-425) Evaluate

$$\frac{\partial}{\partial x} \int_{4}^{6x} \ln(4t) \, \mathrm{d}t$$

a) $\frac{3}{2x}$ b) $6 \ln(24x)$ c) $4 \ln(24x)$ d) $6 \ln(24x) - 24 \ln(2)$ e) $6 \ln(6x) - 24 \ln(2)$ 26) The region bounded by

$y = 3 \sin(x)$

and the *x*-axis, for $0 \le x \le \frac{1}{2} \pi$, is rotated about the line y = -3. The volume of this solid can be represented by:

a)
$$\pi \int_{0}^{\frac{1}{2}\pi} ((3\sin(x) + 3)^{2} - 9) dx$$

b) $2\pi \int_{0}^{\frac{1}{2}\pi} (9\sin(x)^{2} + 3) dx$
c) $\pi \int_{0}^{\frac{1}{2}\pi} (9\sin(x)^{2} - 9) dx$
d) $2\pi \int_{0}^{\frac{1}{2}\pi} 9\sin(x + 3)^{2} dx$
e) $2\pi \int_{0}^{\frac{1}{2}\pi} (3\sin(x) + 3)^{2} dx$

27) Give the third degree Taylor polynomial about x = 1 of $f(x) = \ln(x)$

a) $(x-1) - \frac{1}{3} (x-1)^2 + \frac{1}{5} (x-1)^3$ b) $(x-1) - \frac{1}{2} (x-1)^2 + \frac{1}{6} (x-1)^3$ c) $(x-1) - \frac{1}{2} (x-1)^2 + \frac{1}{3} (x-1)^3$ d) $(x-1) + \frac{1}{3} (x-1)^2 + \frac{1}{3} (x-1)^3$ e) $(x-1) + (x-1)^2 + 2 (x-1)^3$ 28) Which of the following integrals gives the length of the graph of $f(x) = e^{6x}$

for *x* between 0 and 2?

a)
$$\int_{0}^{2} \sqrt{x + e^{12x}} dx$$

b)
$$\int_{0}^{2} \sqrt{1 + e^{12x}} dx$$

c)
$$\int_{0}^{2} \sqrt{x + 36 e^{12x}} dx$$

d)
$$\int_{0}^{2} \sqrt{e^{6x} + 36 e^{12x}} dx$$

e)
$$\int_{0}^{2} \sqrt{1 + 36 e^{12x}} dx$$

29) Find the average value of the function

$$f(x) = e^{5x}$$

over the interval [0, 4].

- a) $\frac{1}{4} (e^{20} 1)$ b) $\frac{1}{5} (e^{20} - 1)$
- c) $\frac{1}{20} e^{20}$ d) $\frac{1}{5} e^{20}$
- e) $\frac{1}{20} (e^{20} 1)$

30) What is the *y*-intercept of the line tangent to the curve $y = x^2 + 8$ at x = 3?

- a) (0,2)
- b) (0,-1)
- c) (0, -10)
- d) (0, 10)
- e) (0, 1)

31) Which of the following function(s) is continuous and differentiable?


- I. $f(x) = \frac{8}{\sqrt{x}}$ II. g(x) = x |x|III. $h(x) = \begin{cases} 7x + 1 & x \le 0 \\ x^2 + 1 & 0 < x \end{cases}$ a) Lonly
- a) I only
- b) II only
- c) III only
- d) I and II only
- e) I and III only
- 32) Find *m*

$$\lim_{x \to 0} \left(\frac{e^{m x^2} - \cos(4 x)}{x^2} \right) = 16$$

a) 4

- b) ₃₂
- c) ₂
- d) 1
- e) 8

33) The graph of the derivative of f is shown below. Which of the following must be true?



a) f has a point of inflection at x = 8. b) f is increasing on [-4, 4]. c) f has a local maximum at x = 0. d) f is concave down on [0, 8]. e) f has a local minimum at x = -4. 34) The sum of two positive integers x and y is 30. Find the value of x that minimizes $P = x^3 - 30 x v$ a) x = 5b) x = 15c) x = 10d) x = 25e) x = 2035) A particle moves on the curve $[x = 6 \sin(t), y = \sin(2t)]$ find the speed of the particle at time $t = \pi$. a) 5.6569 b) 6.3246 c) 6.1644 d) 3.1623 e) 2.8284 36) The function f is defined as $f(x) = \frac{(x-3)^2}{x-7}$ x_{\neq} 7 Which of the following is false? a) f is decreasing on [3, 7].

b) *f* has a vertical asymptote at x = 7.

c) f has a horizontal asymptote at y = 1.

d) *f* has a local maximum at x = 3.

e) *f* is concave up for x > 7.

37) A particle is moving along the *x*-axis and its position at time $t \ge 0$ is given by

$$S(t) = (t-3)^2 (t-6)$$

Which of the following is (are) true?

I. The particle changes direction at x = 3 and x = 6.

II. The particle is slowing down on [0, 3].

- III. The particle is speeding up on [3, 6].
- a) I, II and III
- b) II and III only
- c) I and III only
- d) II only
- e) I only

38) f(x) is a differentiable function and it is decreasing on $(-\infty, \infty)$.

If

$$g(x) = f(12x^3 - x^2)$$

then g has a local maximum at

a) $x = \frac{1}{6}$ b) x = 1c) x = 6d) $x = \frac{1}{18}$ e) x = 0

39) The rate at which a bacteria population grows is proportional to the number of bacteria present. Initially, there were 1000 bacteria present and the population doubled in 6 hours. Roughly how many hours does it take for the population to reach 10000?

- a) 17.4
- b) 31.2
- c) _{13.2}
- d) 19.8
- e) 24.6


40) Find the radius of convergence of the series


$$\sum_{n=1}^{\infty} \frac{n \left(3 x - 9\right)^n}{2^n}$$

a) $\frac{9}{2}$ b) $\frac{2}{9}$

c) $\frac{3}{3}$ d) $\frac{2}{3}$

- e) $\frac{3}{2}$





The graph of g is shown above. Which of the following must be true?

I. $\int_{0}^{6} f(t) dt = 0$ II. $\int_{3}^{6} f(t) dt = 3$ III. $\int_{6}^{0} f(t) dt = -3$ a) II only b) I only c) I and II only d) I and III only e) II and III only 42) If the region bounded by $y = \tan^{-1}(x), y = \frac{1}{4}\pi$ and the y-axis is rotated about the y-axis, the volume of the solid formed is a) 0.413 b) 1.348 c) 0.430

d) 0.674

e) 0.215(x) is represent

43)
$$f(x)$$
 is represented by the Maclaurin series

$$1 - \frac{(2x)^2}{2!} + \frac{(2x)^4}{4!} - \dots + (-1)^n \frac{(2x)^{2n}}{(2n)!} + \dots$$

What is the slope of the line normal to the graph of *f* at $x = \frac{3}{4}\pi$?

- a) _2 b) ₀ c) ₂
- d) $-\frac{1}{2}$

e) $\frac{1}{2}$

44) What are all values of h for which

$$\int_0^\infty \frac{2x}{\left(x^2+1\right)^h} \, \mathrm{d}x$$

converge? a) -1 < h < 1b) $h \ge 1$ c) $h \le 1$ d) h > 1e) h < 145) The base of a solid is the region bounded by

 $y = 5\sqrt{x}$

the x-axis, and

the line x = 5

Each cross-section of the solid perpendicular to the *x*-axis is a square, with one side on the *xy*-plane. Which of the following expressions represents the volume of the solid?

a) $\int_{0}^{5} 25 x \, dx$ b) $\int_{0}^{5} 5 \sqrt{x} \, dx$ c) $\int_{0}^{5} 5 x \, dx$ d) $\int_{0}^{1} 25 x \, dx$ e) $\int_{0}^{1} 5 \sqrt{x} \, dx$ 1) Find

$$\int_{1}^{9} \frac{2}{\sqrt{x}} \, \mathrm{d}x$$

- a) 12 b) 4
- c) 35
- d) 36
- *e) 8
- 2) If

$$f'(x) = -5 (x-6)^2 (x-9)$$

which of the following is true about y = f(x)? a) *f* has a local maximum at x = 6 and a local minimum at x = 9. *b) *f* has a point of inflection at x = 6 and a local maximum at x = 9. c) *f* has a local minimum at x = 6 and a local maximum at x = 9. d) *f* has a point of inflection at x = 6 and a local minimum at x = 9. e) *f* has a local minimum at x = 6 and a point of inflection at x = 9. 3) A curve is described by parametric equations

$$\left[x=5\ln(t),\,y=t^2-4\right]$$

where t > 0. Give an expression for

$$\frac{\partial^2}{\partial x^2}y$$

*a) $\frac{4}{25}t^2$ b) $\frac{4}{5}t$ c) $\frac{4}{25}t$ d) $\frac{2}{5}t^2$ e) $\frac{2}{5}t$

4) Give the value for

$$\lim_{x\to 0} \left(\frac{1}{7} \ \frac{5^x - 1}{x}\right)$$

- a) $7 \ln(5)$
- b) ₍₎
- c) 1

d) The limit does not exist.

$$(1)^{*e} \frac{1}{7} \ln(5)$$

5) Which of the following series converge?

I.

$$\sum_{n=1}^{\infty} \frac{\sin(n)}{n!}$$

II.

$$\sum_{n=1}^{\infty} \frac{x^{2n}}{n!}$$

III.

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)}$$

*a) I, II and III b) II only c) II and III only d) I and II only e) I only 6) If g(f(x)) = x, g(4) = 2 and g'(4) = 10, then f'(2) is a) $\frac{1}{4}$ b) $-\frac{1}{10}$ *c) $\frac{1}{10}$ d) $-\frac{1}{4}$ e) $\frac{2}{5}$ 7) If f is a differentiable function and f(0) = 5 and f(0)

7) If *f* is a differentiable function and f(0) = -5 and f(3) = 10, then which of the following must be true?

I. There exists a *c* in [0,3] where f(c) = 0. II. There exists a *c* in [0,3] where f'(c) = 0. III. There exists a *c* in [0,3] where f'(c) = 5.

a) I only
*b) I and III only
c) II only
d) II and III only
e) I, II and III
8) Evaluate



a)
$$\frac{8}{81}$$

b) $_{0}$
*c) The integral diverges.

d) $\frac{16}{81}$ e) $\frac{4}{81}$

9) Find the area enclosed by the graphs of

$$y = e^{x} + 1$$
$$y = 6$$

and the y-axis.

- a) $6\ln(5) 5$
- b) $6\ln(6) + 6$
- $\frac{c}{6}\ln(5) + 5$
- $^{*d)} 5 \ln(5) 4$
- e) $6\ln(6) 6$
- 10) What is the minimum value of the function

$$f(x) = \frac{2}{\sqrt{x}} + 3\sqrt{x}$$

a) $\frac{2}{3}\sqrt{6}$ b) $\frac{2}{3}$ c) $\frac{1}{3}\sqrt{6}$ *d) $2\sqrt{6}$ e) $\sqrt{6}$ 11) Give the value of

$$\int_{\pi}^{2\pi} \frac{\cos(8x)}{2+\sin(8x)} \, \mathrm{d}x$$

a) $-\frac{1}{8}$ b) $\frac{1}{8}$ c) $\frac{1}{16}$ *d) $_{0}$ e) $_{1}$

12) The side of a cube is expanding at a constant rate of 6 inches per second. What is the rate of change of the surface area, in in^2 per second, when the volume of the cube is 64 in³?

- a) ₃₆
- b) ₁₄₄
- c) ₃₆₀
- *d) ₂₈₈
- e) ₇₂

13) Give the area inside one petal of the polar graph of

 $r = 5 \sin(2 \theta)$

a) $\frac{25}{2} \pi$ b) $\frac{25}{4} \pi$ *c) $\frac{25}{8} \pi$ d) $\frac{25}{16} \pi$ e) $\frac{5}{2}$

14) Give the solution to the initial value problem

$$[y' = 9 x^2 y, y(1) = 1]$$

a) $\frac{1}{3} \ln(x^3) + 1$ b) $e^{3x^3} - 3$ c) $\frac{1}{3} \ln(x^3) + e^{3x^3 - 3}$ e) e^{3x^3}

15) The position of a particle moving along a horizontal line is given by

 $x(t) = 3 (t-4)^3$ What is the maximum speed of the particle for $0 \le t \le 10$?

a) 48 b) 144 c) 576 *d) 324 e) 108 16) $\int [\sec(3x)]^2 dx =$ a) $\frac{1}{3} [\tan(3x)]^2 + C$ b) $-3\tan(3x) + C$ c) $3 [\tan(3x)]^2 + C$ *d) $\frac{1}{3} \tan(3x) + C$ e) $3\tan(3x) + C$ 17) Define the function

 $f(x) = x e^{-8x}$

for x > 0. Give the interval on which the function is increasing. *a) $\begin{pmatrix} 0, \frac{1}{8} \end{pmatrix}$ b) $\begin{pmatrix} 1, 8 \end{pmatrix}$ c) $\begin{pmatrix} 1, \frac{1}{8} \\ e \end{pmatrix}$

$$\begin{array}{c} d) \left(0, \frac{1}{8} e\right) \\ e) (0, 8) \\ 18) \\ \hline \\ & & & & \\ & & & & \\ & & & \\ & & &$$

Which of the following differential equations correspond to the slope field shown in the figure above?

*a) $\frac{dy}{dx} = -\frac{x}{y}$ b) $\frac{dy}{dx} = \frac{y}{x}$ c) $\frac{dy}{dx} = -\frac{y}{x}$ d) $\frac{dy}{dx} = \frac{x}{y}$ e) $\frac{dy}{dx} = 5 x y$ 19) Evaluate $\lim_{h \to 0} \left(\frac{\cos(5x+5h) - \cos(5x)}{h} \right)$ $(a)^{*a} - 5\sin(5x)$ b) $5 \sin(5x)$ c) $5\cos(5x)$ d) $-5\cos(5x)$ e) The limit does not exist. 20) If $\int_{0}^{16} e^{x} dx = m$ then $\int_{0}^{4} x e^{x^{2}} dx$ is *a) $\frac{1}{2}m$ b) _m c) 2 m d) _m2 e) $\frac{1}{2}m^2$ 21) Find the area of the region enclosed by the graph of

and the line

$$y=2x^2$$

$$y = 3x$$

*a) $\frac{9}{8}$ b) $\frac{9}{4}$ c) $\frac{3}{8}$ d) $\frac{4}{27}$ e) $\frac{8}{27}$ 22) Suppose

 $z = e^{y}$ $y = 4x^{3} - 4$ $x = 1 + Ln(t^{2})$

and

- What is dz/dt when t = 1?
- a) 6
- b) 4
- c) ₁₂
- d) ₁₆
- *e) 24
- 23) Evaluate

$$\int_{0}^{\frac{1}{2}\pi} \cos^2\left(13x\right) \sin(13x) \, \mathrm{d}x$$

a) $-\frac{1}{78}$ b) $\frac{1}{78}$ c) $-\frac{1}{39}$ *d) $\frac{1}{39}$ e) $\frac{1}{13}$

24) Give an equation for the tangent line to the parametric curve $[x = e^t, y = t^2 + 4t]$

at t = 0. a) $y = \frac{4(x-1)}{e}$ *b) y = 4x-4c) y = 4e(x-1) d) y - 1 = 4xe) y - 1 = 4x - 425) Evaluate

$$\frac{\partial}{\partial x} \int_{4}^{6x} \ln(4t) \, \mathrm{d}t$$

a) $\frac{3}{2x}$ *b) $6 \ln(24 x)$ c) $4 \ln(24x)$ d) $6\ln(24x) - 24\ln(2)$ e) $6\ln(6x) - 24\ln(2)$ 26) The region bounded by

$$y = 3 \sin(x)$$

 $y = 3 \operatorname{sm}(x)$ and the *x*-axis, for $0 \le x \le \frac{1}{2} \pi$, is rotated about the line y = -3. The volume of this solid can be represented by:

*a)
$$\pi \int_{0}^{\frac{1}{2}\pi} ((3\sin(x) + 3)^{2} - 9) dx$$

b) $2\pi \int_{0}^{\frac{1}{2}\pi} (9\sin(x)^{2} + 3) dx$
c) $\pi \int_{0}^{\frac{1}{2}\pi} (9\sin(x)^{2} - 9) dx$
d) $2\pi \int_{0}^{\frac{1}{2}\pi} 9\sin(x + 3)^{2} dx$
e) $2\pi \int_{0}^{\frac{1}{2}\pi} (3\sin(x) + 3)^{2} dx$

27) Give the third degree Taylor polynomial about x = 1 of $f(x) = \ln(x)$

a)
$$(x-1) - \frac{1}{3} (x-1)^2 + \frac{1}{5} (x-1)^3$$

b) $(x-1) - \frac{1}{2} (x-1)^2 + \frac{1}{6} (x-1)^3$
*c) $(x-1) - \frac{1}{2} (x-1)^2 + \frac{1}{3} (x-1)^3$
d) $(x-1) + \frac{1}{3} (x-1)^2 + \frac{1}{3} (x-1)^3$
e) $(x-1) + (x-1)^2 + 2 (x-1)^3$
28) Which of the following integrals gives the length of the graph of $f(x) = e^{6x}$

for x between 0 and 2?

a)
$$\int_{0}^{2} \sqrt{x + e^{12x}} dx$$

b)
$$\int_{0}^{2} \sqrt{1 + e^{12x}} dx$$

c)
$$\int_{0}^{2} \sqrt{x + 36 e^{12x}} dx$$

d)
$$\int_{0}^{2} \sqrt{e^{6x} + 36 e^{12x}} dx$$

*e)
$$\int_{0}^{2} \sqrt{1 + 36 e^{12x}} dx$$

29) Find the average value of the function

$$f(x) = e^{5x}$$

over the interval [0, 4].

- a) $\frac{1}{4} (e^{20} 1)$ b) $\frac{1}{5} (e^{20} - 1)$
- c) $\frac{1}{20} e^{20}$ d) $\frac{1}{5} e^{20}$ *e) $\frac{1}{20} (e^{20} - 1)$

30) What is the *y*-intercept of the line tangent to the curve $y = x^2 + 8$ at x = 3?

- a) (0, 2)
- *b) (0, -1)
- c) (0, -10)
- d) (0, 10)
- e) (0, 1)

31) Which of the following function(s) is continuous and differentiable?


- I. $f(x) = \frac{8}{\sqrt{x}}$ II. g(x) = x |x|III. $h(x) = \begin{cases} 7x + 1 & x \le 0 \\ x^2 + 1 & 0 < x \end{cases}$ *a) I only b) II only
- c) III only
- d) I and II only
- e) I and III only
- 32) Find *m*

$$\lim_{x \to 0} \left(\frac{e^{m x^2} - \cos(4 x)}{x^2} \right) = 16$$

a) 4

- b) ₃₂
- c) 2
- d) 1
- *e) 8

33) The graph of the derivative of f is shown below. Which of the following must be true?



*a) f has a point of inflection at x = 8. b) f is increasing on [-4, 4]. c) f has a local maximum at x = 0. d) f is concave down on [0, 8]. e) f has a local minimum at x = -4. 34) The sum of two positive integers x and y is 30. Find the value of x that minimizes $P = x^3 - 30 x v$ a) x = 5b) x = 15 $(*c)_{x=10}$ d) x = 25e) x = 2035) A particle moves on the curve $[x = 6 \sin(t), y = \sin(2t)]$ find the speed of the particle at time $t = \pi$. a) 5.6569 *b) 6.3246 c) 6.1644 d) 3.1623 e) 2.8284 36) The function f is defined as $f(x) = \frac{(x-3)^2}{x-7}$ x_{\neq} 7 Which of the following is false?

which of the following is laise

a) f is decreasing on [3, 7].

b) *f* has a vertical asymptote at x = 7.

*c) *f* has a horizontal asymptote at y = 1.

d) *f* has a local maximum at x = 3.

e) *f* is concave up for x > 7.

37) A particle is moving along the *x*-axis and its position at time $t \ge 0$ is given by

$$\mathcal{S}(t) = (t-3)^2 (t-6)$$

Which of the following is (are) true?

I. The particle changes direction at x = 3 and x = 6.

II. The particle is slowing down on [0, 3]. III. The particle is speeding up on [3, 6]. a) I, II and III b) II and III only c) I and III only *d) II only e) I only 38) f(x) is a differentiable function and it is decreasing on $(-\infty, \infty)$. If

$$g(x) = f\left(12 x^3 - x^2\right)$$

then g has a local maximum at

a) $x = \frac{1}{6}$ b) x = 1c) x = 6*d) $x = \frac{1}{18}$ e) x = 0

39) The rate at which a bacteria population grows is proportional to the number of bacteria present. Initially, there were 1000 bacteria present and the population doubled in 6 hours. Roughly how many hours does it take for the population to reach 10000?

a) 17.4


- b) 31.2
- c) _{13.2}
- *d) _{19.8}
- e) 24.6


40) Find the radius of convergence of the series

$$\sum_{n=1}^{\infty} \frac{n \left(3 x - 9\right)^n}{2^n}$$

- a) $\frac{9}{2}$ b) $\frac{2}{9}$

- c) $\frac{1}{3}$ *d) $\frac{2}{3}$
- e) $\frac{3}{2}$





The graph of g is shown above. Which of the following must be true?

I. $\int_{0}^{6} f(t) dt = 0$ II. $\int_{3}^{6} f(t) dt = 3$ III. $\int_{6}^{0} f(t) dt = -3$ a) II only b) I only *c) I and II only d) I and III only e) II and III only 42) If the region bounded by $y = \tan^{-1}(x), y = \frac{1}{4}\pi$ and the y-axis is rotated about the y-axis, the volume of the solid formed is a) 0.413 b) 1.348 c) 0.430 *d) 0.674

e) 0.215

43) f(x) is represented by the Maclaurin series

$$1 - \frac{(2x)^2}{2!} + \frac{(2x)^4}{4!} - \dots + (-1)^n \frac{(2x)^{2n}}{(2n)!} + \dots$$

What is the slope of the line normal to the graph of *f* at $x = \frac{3}{4}\pi$?

a) $_{-2}$ b) $_{0}$ c) $_{2}$ *d) $_{-\frac{1}{2}}$ e) $\frac{1}{2}$

44) What are all values of h for which

$$\int_0^\infty \frac{2x}{\left(x^2+1\right)^h} \, \mathrm{d}x$$

converge? a) -1 < h < 1b) $h \ge 1$ c) $h \le 1$ *d) h > 1e) h < 145) The base of a solid is the region bounded by

 $y = 5\sqrt{x}$

the x-axis, and

the line x = 5

Each cross-section of the solid perpendicular to the *x*-axis is a square, with one side on the *xy*-plane. Which of the following expressions represents the volume of the solid?

*a) $\int_{0}^{5} 25 x \, dx$ b) $\int_{0}^{5} 5 \sqrt{x} \, dx$ c) $\int_{0}^{5} 5 x \, dx$ d) $\int_{0}^{1} 25 x \, dx$ e) $\int_{0}^{1} 5 \sqrt{x} \, dx$